

## **Short-term responses of greenhouse gas emissions and ecosystem carbon fluxes to elevated ozone and N fertilization in a temperate grassland**

Wang, Jinyang; Hayes, Felicity; Hill, Paul W.; Chadwick, David R.; Mills, Gina; Jones, Davey L.

### **Atmospheric Environment**

DOI:

[10.1016/j.atmosenv.2019.05.027](https://doi.org/10.1016/j.atmosenv.2019.05.027)

Published: 15/08/2019

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Wang, J., Hayes, F., Hill, P. W., Chadwick, D. R., Mills, G., & Jones, D. L. (2019). Short-term responses of greenhouse gas emissions and ecosystem carbon fluxes to elevated ozone and N fertilization in a temperate grassland. *Atmospheric Environment*, 211, 204-213. <https://doi.org/10.1016/j.atmosenv.2019.05.027>

#### **Hawliau Cyffredinol / General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1    **Short-term responses of greenhouse gas emissions and ecosystem carbon fluxes to**  
2    **elevated ozone and N fertilization in a temperate grassland**

3    Jinyang Wang<sup>a,\*</sup>, Felicity Hayes<sup>b</sup>, David R. Chadwick<sup>a</sup>, Paul W. Hill<sup>a</sup>, Gina Mills<sup>b</sup>, Davey L.  
4    Jones<sup>a,c</sup>

5    <sup>a</sup> *School of Natural Sciences, Environment Centre Wales, Bangor University, Bangor, Gwynedd*  
6    *LL57 2UW, UK*

7    <sup>b</sup> *Centre for Ecology and Hydrology, Environment Centre Wales, Bangor, Gwynedd LL57 2UW,*  
8    *UK*

9    <sup>c</sup> *UWA School of Agriculture and Environment, University of Western Australia, Crawley, WA*  
10    *6009, Australia*

11    \*Corresponding author.

12    E-mail address: jy.wang@bangor.ac.uk (J. Wang)

13

14

15

16

## Abstract

Growing evidence suggests that tropospheric ozone has widespread effects on vegetation, which can contribute to alter ecosystem carbon (C) dynamics and belowground processes. In this study, we used intact soil mesocosms from a semi-improved grassland and investigated the effects of elevated ozone, alone and in combination with nitrogen (N) fertilization on soil-borne greenhouse gas emissions and ecosystem C fluxes. Ozone exposure under fully open-air field conditions was occurred during the growing season. Across a one-year period, soil methane ( $\text{CH}_4$ ) and nitrous oxide ( $\text{N}_2\text{O}$ ) emissions did not differ between treatments, but elevated ozone significantly depressed soil  $\text{CH}_4$  uptake by 14% during the growing season irrespective of N fertilization. Elevated ozone resulted in a 15% reduction of net ecosystem exchange of carbon dioxide, while N fertilization significantly increased ecosystem respiration during the growing season. Aboveground biomass was unaffected by elevated ozone during the growing season but significantly decreased by 17% during the non-growing season. At the end of the experiment, soil mineral N content, net N mineralization and extracellular enzyme activities (i.e., cellobiohydrolase and leucine aminopeptidase) were higher under elevated ozone than ambient ozone. The short-term effect of single application of N fertilizer was primarily responsible for the lack of the interaction between elevated ozone and N fertilization. Therefore, results of our short-term study suggest that ozone exposure may have negative impacts on soil  $\text{CH}_4$  uptake and C sequestration and contribute to accelerated rates of soil N-cycling.

37    *Keywords:* Air pollutant; Fertilizer management; Ground level O<sub>3</sub>; Plant-soil feedbacks; Soil

38    nutrient cycling

39

## 1. Introduction

Owing to the global increase in ozone precursor emissions (i.e., nitrogen oxides, carbon monoxide and volatile organic compounds), further increases of background ozone concentrations in the Northern Hemisphere may occur over this century unless precursor emissions are effectively controlled (Fowler et al., 2008; Meehl et al., 2007). Tropospheric ozone is not only the third-most-important contributor to the human-induced greenhouse gas (GHG) effect after carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) (IPCC, 2007), but also the important gaseous air pollutant in terms of its effects on net primary production, soil carbon (C) sequestration, as well as other ecosystem services (Ainsworth et al., 2012; Sicard et al., 2017).

Ozone may have important indirect effects on global change through its effects on CH<sub>4</sub> and nitrous oxide (N<sub>2</sub>O) emissions. In the past decade, the effects of elevated ozone on CH<sub>4</sub> and N<sub>2</sub>O emissions, especially from agroecosystems, have been increasingly studied. For example, studies in rice fields conducted under either free-air open conditions or the open-top chambers (OTC) have consistently shown that CH<sub>4</sub> emissions were substantially less under elevated ozone than under ambient or charcoal-filtered air conditions (Bhatia et al., 2011; Tang et al., 2015; Zheng et al., 2011). The impacts of elevated ozone on N<sub>2</sub>O emissions have been examined in a range of agricultural systems, including rice fields (Bhatia et al., 2011; Kou et al., 2015; Tang et al., 2015), a soybean field (Decock et al., 2012), an annual grassland (Sánchez-Martín et al., 2017) and a meadow ecosystem (Kanerva et al., 2007). However, the results from these experiments are conflicting, showing either positive (Sánchez-Martín et al.,

2017), negative (Bhatia et al., 2011; Kanerva et al., 2007) or no significant (Decock et al., 2012; Kou et al., 2015; Tang et al., 2015) responses of N<sub>2</sub>O emissions to ozone exposure. Such inconsistencies may stem from the complex impacts of other factors (e.g. exposure methodology, exposure duration and plant type etc.) on the responses of N<sub>2</sub>O emissions to elevated ozone (Tang et al., 2015).

On the other hand, ozone may also have important indirect effects on global change through its effect on terrestrial C sequestration. At the global scale, Sitch et al. (2007) used a global land C cycle model by including the ozone deposition effect and demonstrated that the negative impact of elevated ozone on plant productivity may result in a significant suppression of the global land-C sink. Such modeling studies (Sitch et al., 2007), however, are based on the effects of ozone on photosynthetic rates and aboveground growth, and do not consider its effects on soil C fluxes. Furthermore, studies now support the view that ozone may also have a profound impact on belowground processes because of altered C allocation to the roots and associated rhizosphere (Andersen, 2003; Fuhrer et al., 2016; Grantz et al., 2006; Wang et al., 2019). To better understand how increasing tropospheric ozone will affect global C fluxes, studies addressing ozone effects on soil C fluxes are therefore required.

Net ecosystem exchange (NEE) of CO<sub>2</sub> represents the balance between gross ecosystem productivity (GEP) and ecosystem respiration ( $R_{eco}$ ) (Niu et al., 2010). It has been suggested that directly measuring NEE can be used to evaluate the response of ecosystem C sequestration to climatic factors (Luyssaert et al., 2007). To date, divergent responses of ecosystem C fluxes

to elevated ozone have been reported in an annual grassland (Calvete-Sogo et al., 2014), peatlands (Haapala et al., 2011; Toet et al., 2011), and a subalpine grassland (Volk et al., 2011). These contrasting results of studies from different ecosystem types under distinct ozone exposure conditions signify a limited understanding of ozone effects on ecosystem C fluxes.

Despite considerable awareness of the potential interaction between elevated ozone and nitrogen (N) fertilization, very little attention has been paid to investigate their combined effects on ecosystem production and function (Mills et al., 2016). In a seven-year study from a subalpine grassland, plant biomass was positively affected by N addition but neither ozone nor its interaction with N addition (Volk et al., 2014). In contrast, results of a recent meta-analysis show that elevated ozone generally tends to exacerbate the negative impact of limiting N on plant growth (Yendrek et al., 2013). Similarly, in an annual grassland ozone-induced reductions in N content and C/N ratio of plant biomass can be counterbalanced by N addition (Sánchez-Martín et al., 2017), despite the absence of their interaction on GHG emissions and ecosystem C fluxes. Altogether, these conflicting results suggest that the direction and magnitude of the combined effect of elevated ozone and N fertilization on ecosystem processes remains unpredictable.

As the dominant form of agriculture by land area, grasslands represent over two thirds of utilized agricultural land area in the UK (Defra, 2016). Semi-natural vegetation is well known to be adversely affected by elevated ozone concentrations across Europe (Hayes et al., 2007; Mills et al., 2011). The aforementioned studies have mainly focused on croplands and annual

grasslands, but less is known about how elevated ozone and its interaction with N fertilization affect ecosystem production and function (e.g. fodder production, trace gas emissions and soil properties) in temperate grasslands (Wang et al., 2019). Our objectives for this study were therefore (i) to determine the effects of elevated ozone on CH<sub>4</sub> and N<sub>2</sub>O fluxes from a temperate grassland in the UK throughout a one-year period; (ii) to investigate how elevated ozone would affect ecosystem C fluxes during the growing season (i.e., ozone exposure period); and (iii) to test whether the responses of these fluxes to elevated ozone would be interacted with N fertilization. To this end, we carried out a field experiment with intact soil mesocosms from a temperate semi-improved grassland and exposed them to elevated ozone for one growing season using a ozone free-air controlled enrichment (O<sub>3</sub>-FACE) platform.

## **2. Materials and methods**

### *2.1 Experimental design*

The experiment was conducted in 2017 in a O<sub>3</sub>-FACE system at CEH Bangor Air Pollution Facility, Abergwyngregyn, North Wales, UK (13 m asl, 53°15'N, 4°01'W). The climate at the site is classed as temperate-oceanic with a mean annual soil temperature of 11°C at 10 cm depth and a mean annual rainfall of 1250 mm. Experimental plots consisted of 16 intact soil mesocosms (31 cm diameter × 25 cm deep), which were excavated in the early spring of 2017 from a semi-improved upland grassland located at the Henfaes Research Station, Abergwyngregyn, North Wales, UK (53°13'N, 4°0'W). The upland site is located at approximately 270 m altitude with a mean growing season temperature of 10 °C and a mean



annual precipitation of 1200 mm. This semi-improved grassland site had not received any fertilizer applications and is grazed at low stocking densities (1-2 ewes ha<sup>-1</sup>). The vegetation was classified as *Cynosurus cristatus*-*Centaurea nigra* grassland (NVC MG5; Rodwell, 1992). The soil is classified as an Orthic Podzol (FAO, 1981). The soil of this site had a slightly acidic pH of 5.39±0.04, total soil C of 8.45±0.09%, and total N of 0.81±0.01%.

Four treatments were established to determine the effects of elevated ozone, N fertilization and their potential interaction. The low and high ozone treatments from the O<sub>3</sub>-FACE system as described below (hereafter called the ambient and elevated ozone treatments, respectively) were used in this study. The exposure period of ozone was from May 26 to October 9, 2017. Each quarter of these mesocosms was randomly assigned to one of the four combinations of ozone and N fertilizer treatments. Nitrogen was added as NH<sub>4</sub>NO<sub>3</sub> at a rate of 100 kg N ha<sup>-1</sup> dissolved in 200 mL of deionized water per mesocosm on May 26, 2017. For the control treatment, an equal amount of deionized water was added. To ensure all other macro-nutrients were not limiting pasture growth, and that any nutrient response was the result of the N addition, calcium superphosphate for P and potassium chloride for K were also used and applied to all treatments at a rate of 10 and 50 kg ha<sup>-1</sup>, respectively. The fertilizer addition rates were based on national guidelines.

The O<sub>3</sub>-FACE system was established in the spring of 2014, consisting of nine rings of 4 m diameter as described previously (Wang et al., 2019). Briefly, the rings were arranged in a replicated 3 × 3 Latin square with 10 m between the centers of each ring. Low (ambient air)

and high ozone (ambient air + 20 ppb) treatments were used in this study. Ozone was generated from oxygen concentrated from air (Integra 10, SeQual) using a G11 ozone generator (Pacific Ozone). Small fans (200 mm, Explair) were used to push the ozone through the delivery pipe (65 mm, with 3 mm holes every 10 cm). Ozone delivery was achieved via computer controlled (LabView version 2012) solenoid valves operating using pulse width modulation. Wind speed was monitored continuously (WindSonic, Gill Instruments Ltd, UK) and was used to instantaneously adjust solenoid operation and thus ozone delivery. Ozone release was reduced at wind speeds below  $16 \text{ m s}^{-1}$  and did not occur when wind speeds fell below  $2 \text{ m s}^{-1}$ . At very high wind speeds the ozone concentrations may not be well controlled and thus did not reach the target maximum concentrations. Despite this, we still got elevated ozone with the ‘high’ ozone treatment compared to the low ozone treatment as the solenoid valves were  $<1 \text{ m}$  from the  $\text{O}_3$ -FACE rings, the response time of ozone delivery to track windspeed was fast. Ozone was sampled adjacent to the plants in each ring at a height of 30 cm for approximately 3.5 min in every 30-min using an ozone analyzer (ThermoScientific, Model 49i, Reading, UK).

## *2.2 Soil GHG emissions and ecosystem C fluxes*

Soil GHG fluxes were measured throughout the experimental year. Gas samples were collected by placing a handmade opaque chamber over the intact soil mesocosms and sealing with a wide rubber band to ensure that the headspace inside the chamber was air-tight. Three gas samples were taken within a 40-min enclosure time and immediately transferred into pre-evacuated 20-mL screw-cap vials (QUMA Elektronik & Analytik GmbH, Germany). Gas samples were

stored under positive pressure and analyzed using a Perkin Elmer Clarus 580 gas chromatograph equipped with a Turbo Matrix 110 autosampler (PerkinElmer Inc., Shelton, CT, USA). CH<sub>4</sub> and CO<sub>2</sub> were detected with a flame ionization detector connected to a methanizer and N<sub>2</sub>O with a <sup>63</sup>Ni electron-capture detector. Three standard gases were used for calibration with concentrations of 1.42, 2.92, and 10.4 ppm CH<sub>4</sub>; 258, 496 and 1276 ppm CO<sub>2</sub>; and 308, 641, and 1500 ppb N<sub>2</sub>O (BOC Gases Ltd, Guildford, UK). Gas fluxes were calculated from the concentration change in the chamber versus time and were adjusted for air temperature and atmospheric pressure at the time of sampling. The annual fluxes of these gases were approximated by applying the trapezoid rule assuming constant flux rates per day.

Ecosystem C fluxes were measured during the growing season using a static-chamber method, which has been used and validated in previous studies (Niu et al., 2010; Volk et al., 2011). The measurements were made from June to September in 2017 and was taken between 8:00 and 11:00 am on sunny days. If it is rained or cloudy on a scheduled sampling date, we postponed the measurements and selected the next sunny day. In total, complete sets of measurements were made seven times during the growing season. A portable infrared gas analyzer (EGM-4 Environmental Gas Monitor for CO<sub>2</sub>; PP Systems Ltd, Hitchin, UK), which was attached to a handmade transparent plastic chamber, was employed to sample and measure CO<sub>2</sub> concentration ([CO<sub>2</sub>]) *in situ* as described elsewhere (Williamson et al., 2016). A small fan created moderate turbulence inside the chamber to facilitate air mixing. A temperature probe was also installed inside to determine the chamber air temperature. During the measurement, the chamber was tightly sealed to the rim of the mesocosm using a cell foam band. The chamber

[CO<sub>2</sub>] were consecutively recorded at 20 s intervals during a 2-min measurement period per mesocosm. The chamber [CO<sub>2</sub>] did not drop below 340 ppm or rise above 500 ppm. After the net ecosystem exchange (NEE) of CO<sub>2</sub> measurement, the chambers were opened to allow the chamber [CO<sub>2</sub>] to return to the ambient atmospheric concentration. Subsequently, the chamber was again placed over the mesocosm and covered with a shade cloth, and the [CO<sub>2</sub>] measurement was repeated over a 2-min period (i.e., dark respiration). Due to the light removal inside the chamber, the data of the chamber [CO<sub>2</sub>] were considered to represent ecosystem respiration ( $R_{eco}$ ). To eliminate the disturbance effect during measurement, the first 20 s of data was omitted in subsequent analysis to allow for initial adjustment of the chamber [CO<sub>2</sub>]. The quality of the measurement was considered acceptable if a linear regression of [CO<sub>2</sub>] versus time during the following 100 s yielded  $R^2 \geq 0.9$ , indicating strictly linear changes in the chamber [CO<sub>2</sub>]. Gross ecosystem productivity (GEP) was computed by the sum of absolute values of NEE and  $R_{eco}$ . By convention, negative NEE values refer net C uptake by the ecosystem, while positive NEE values represent net C loss from the ecosystem.

### *2.3 Soil abiotic and biotic properties and plant analysis*

Air temperature and rainfall were recorded hourly on site with an automated weather station. Parallel to gas sampling and ecosystem C flux measurements, soil temperature and volumetric water content (v/v, %) at a depth of 10 cm in each mesocosm were measured by a handheld thermometer (HI98509 Checktemp®1; Hanna Instruments Ltd, Leighton Buzzard, UK) and moisture sensor (ML3 ThetaProbe; Delta-T, Cambridge, UK), respectively. Water-filled pore

space (WFPS) was calculated by dividing volumetric water content by total porosity (Linn and Doran, 1984). Total porosity was calculated as  $[1 - (\text{bulk density} / \text{particle density})] \times 100\%$  and using a particle density of  $2.65 \text{ g cm}^{-3}$ .

In May 2018, two soil cores (10-cm deep; 5-cm diameter) were collected from each mesocosm. Fresh soils were passed through a 2-mm sieve to remove visible plant material and small stones, and then placed at  $4^\circ\text{C}$  to await analysis. Soil water content was determined gravimetrically by drying soil in an oven (24 h,  $105^\circ\text{C}$ ). Available soil C and N pools were quantified by extracting soil subsamples with  $0.5 \text{ M K}_2\text{SO}_4$  (1:5 w/v). The concentrations of ammonium ( $\text{NH}_4^+$ ) and nitrate ( $\text{NO}_3^-$ ) in the extracts were determined via the colorimetric salicylate procedure of Mulvaney (1996) and the vanadate method of Miranda et al. (2001), respectively. For soil microbial biomass, additional subsamples were fumigated for 24 h with chloroform and similarly extracted with  $0.5 \text{ M K}_2\text{SO}_4$  (1:5 w/v) (Vance et al., 1987). Dissolved organic C (DOC) and total dissolved N (TDN) in the  $0.5 \text{ M K}_2\text{SO}_4$  extracts (fumigation and non-fumigation samples) were quantified using a Multi N/C 2100 TOC analyzer (AnalytikJena, Jena, Germany). Dissolved organic N (DON) was calculated as the difference between TDN and soil mineral N. Microbial biomass C and N concentrations were corrected using correction factors of 0.45 for C and 0.54 for N (Brookes et al., 1985; Wu et al., 1990). Total C (TC) and N (TN) contents of ground soil and plant samples were determined with a TruSpec<sup>®</sup> elemental analyzer (Leco Corp., St Joseph, MI). Net N mineralization and nitrification rates were determined by the aerobic incubation of soil samples for 14 days at  $10^\circ\text{C}$  in the dark (Hart et al., 1994), followed by extraction with  $0.5 \text{ M K}_2\text{SO}_4$  and analyzing for soil mineral N as

described above. To assess the effects of elevated ozone and N fertilization on hydrolytic extracellular enzyme activities, the potential activities of four extracellular enzymes:  $\beta$ -glucosidase, cellobiohydrolase, N-acetyl-glucosaminidase and leucine aminopeptidase were measured according to the fluorometric protocol (DeForest, 2009; Saiya-Cork et al., 2002).


Plant biomass was measured twice by cutting all plants at 2 cm above the ground at the end of growing season (early November of 2017) and in May 2018, respectively. For root biomass measurement, two additional soil cores (10-cm deep; 5-cm diameter) was sampled from each mesocosm. The harvested soil cores were rinsed thoroughly and passed through a 0.5-mm aperture sieve and the root fragments remaining in the sieve recovered. All plant samples were oven-dried at 65 °C and weighed to determine their dry biomass.

#### *2.4 Statistical analysis*

Each parameter was tested for normal distribution (using Shapiro-Wilk's test) and equality of variance (using Levene's test), and parameters with non-normal distributions or unequal variances were either logarithmically or square-root transformed when necessary. For analysis of time-series data (i.e., repeated measurements of soil gas fluxes and soil factors), we used linear mixed effects models (LME, *package* LME4; Bates et al. 2014) to test for the fixed effects of ozone, N fertilization and their possible interaction. The spatial replication and time (sampling days) were included as random effects. If the Akaike's information criterion (AIC) showed an improvement in the LME models, we included a first-order temporal autoregressive function to account for the decreasing correlation of the measurements with increasing time

242 and/or a variance function (varIdent) to account for heteroscedasticity in the fixed-factor  
243 variances (Crawley, 2012). We used linear mixed effects models to test the fixed effects of  
244 ozone, N fertilization and their interaction on soil properties, extracellular enzyme activities  
245 and plant biomass. Multiple comparisons were made using the Tukey HSD test.

246 To assess the relationships between soil GHG emissions or ecosystem C fluxes and soil  
247 factors (temperature and WFPS), we used the mean values of the four replicate mesocosms on  
248 each sampling day, and conducted Pearson correlation tests over the entire sampling period. In  
249 all statistical tests, differences among treatments were considered significant at  $P \leq 0.05$  and  
250 marginally significant at  $P \leq 0.09$ . Statistical analyses were performed in R version 3.2.2 (R  
251 Development Core Team, 2015).



**Figure 1.** Daily maximum and minimum air temperature and daily rainfall (A) over the period of May 25, 2017 to May 31, 2018, daily ozone concentration (B) and AOT40 during daylight hours (C) from ambient and elevated ozone treatments in the 2017 growing season. Values represent mean $\pm$ SEM ( $n = 3$ ).

### 3. Results

#### 3.1 Climatic conditions and ozone exposure


Over the experimental period, air temperature ranged from  $-5^{\circ}\text{C}$  to  $30^{\circ}\text{C}$ , with an annual average of  $11^{\circ}\text{C}$  (Fig. 1A). The total rainfall was 324 mm, of which only 13% occurred during



the growing season (May–September 2017). Intact soil mesocosms were exposed to ozone in the field from May 25 to October 9, 2017, with a total of 136 days effective exposure (Fig. 1B). During the exposure period, the 24-h mean ozone concentrations were  $22.9 \pm 0.6$  ppb and  $54.9 \pm 6.1$  ppb in the ambient and elevated ozone treatments, respectively. As shown in Fig. 1C, AOT40 (accumulated ozone exposure over a threshold of 40 ppb) in the elevated ozone treatment was  $31.9 \pm 10.1$  ppm.h being markedly higher than that ( $< 0.5$  ppm.h) in the ambient treatment.

### *3.2 Seasonal variability of soil microclimate and GHG emissions*

Soil temperature followed a clear seasonal pattern with an annual mean of  $13.7$  °C across all treatments (Fig. 2A). Soil temperatures were highest in July 2017 and lowest during January 2018. Soil WFPS varied seasonally in response to rainfall and ranged from 12.6% to 87.6% (Fig. 2B). Soil WFPS periodically increased following summer rainfall. Soil water contents steadily increased during the winter season when a large proportion of the annual rainfall occurred. Neither soil temperature nor WFPS differed between treatments (all  $P > 0.1$ ).



**Figure 2.** Soil temperature (A), water-filled pore space (WFPS) (B) at a depth of 10 cm, and year-round fluxes of  $\text{CH}_4$  (C),  $\text{N}_2\text{O}$  (D) and night-time  $\text{CO}_2$  (E) from a temperate semi-improved grassland exposed to different levels of ozone and N. The averages of soil microclimate parameters and gas fluxes over the experimental year are shown in the right panel. Values represent  $\text{mean} \pm \text{SEM}$  ( $n = 4$ ). The horizontal dashed line marks the zero flux. Grey shading

marks the ozone exposure period during the growing season. Results from the linear mixed effects model at  $P \leq 0.05$  are shown.

This semi-improved grassland was a small sink of atmospheric CH<sub>4</sub>. Across all treatments, CH<sub>4</sub> fluxes ranged from -84.1 to 23.9  $\mu\text{g C m}^{-2} \text{ h}^{-1}$  (Fig. 2C). During the phase of ozone exposure, CH<sub>4</sub> uptake rates were significantly affected by elevated ozone with a 14% reduction ( $P = 0.043$ ) regardless of N fertilizer. A marginally significant interaction between elevated ozone and N fertilizer was detected during the winter season ( $P = 0.077$ ). In parallel with high soil WFPS, CH<sub>4</sub> fluxes tended to be high and occasionally positive during the winter season. Overall, CH<sub>4</sub> fluxes were negatively correlated with soil temperature and positively correlated with soil WFPS ( $r = -0.41$  and  $0.51$ , both  $P < 0.001$ ; Table 1), indicative of an increase in CH<sub>4</sub> uptake rates with an increase in soil temperature but a decrease of that with an increase in soil water content. Because of the large spatial and temporal variations, neither elevated ozone nor N fertilization had a significant effect on CH<sub>4</sub> emissions throughout the experimental period.

As with CH<sub>4</sub>, N<sub>2</sub>O emissions were temporally and spatially variable (Fig. 2D). Across all treatments, N<sub>2</sub>O fluxes ranged from -11.6 to 33.3  $\mu\text{g N m}^{-2} \text{ h}^{-1}$ . On a few occasions, fluxes were negative indicating the occurrence of N<sub>2</sub>O consumption by the soil. In contrast to CH<sub>4</sub>, N<sub>2</sub>O fluxes correlated positively with soil temperature and negatively with soil WFPS ( $r = 0.36$  and  $-0.33$ , both  $P < 0.001$ ; Table 1). For both seasons, neither elevated ozone nor N fertilizer affected N<sub>2</sub>O emissions.

**Table 1** Pearson correlation coefficients between greenhouse gas (CH<sub>4</sub>, N<sub>2</sub>O and night-time CO<sub>2</sub>) fluxes or ecosystem C fluxes and soil environmental factors (soil temperature and water-filled pore space (WFPS) at a depth of 10 cm). NEE, net ecosystem exchange of CO<sub>2</sub>;  $R_{eco}$ , ecosystem respiration; GEP, gross ecosystem productivity.


| Parameter                    | <i>n</i> | Soil temperature     | WFPS                 |
|------------------------------|----------|----------------------|----------------------|
| <i>Greenhouse gas fluxes</i> |          |                      |                      |
| CH <sub>4</sub>              | 123      | -0.41 <sup>***</sup> | 0.51 <sup>***</sup>  |
| N <sub>2</sub> O             | 123      | 0.36 <sup>***</sup>  | -0.33 <sup>***</sup> |
| Night-time CO <sub>2</sub>   | 123      | 0.69 <sup>***</sup>  | -0.53 <sup>***</sup> |
| <i>Ecosystem C fluxes</i>    |          |                      |                      |
| NEE                          | 28       | 0.04                 | -0.67 <sup>***</sup> |
| $R_{eco}$                    | 28       | 0.45 <sup>*</sup>    | 0.44 <sup>*</sup>    |
| GEP                          | 28       | 0.26                 | 0.67 <sup>***</sup>  |

<sup>\*</sup>, <sup>\*\*</sup>, <sup>\*\*\*</sup> Values are significantly different from zero at  $P < 0.05$ , 0.01, and 0.001, respectively

The variation in seasonal night-time CO<sub>2</sub> fluxes was more pronounced as compared with CH<sub>4</sub> and N<sub>2</sub>O emissions (Fig. 2E). Greatest fluxes occurred during the growing season despite a few occasions when small fluctuations occurred. Lowest emissions occurred after the autumn harvest of aboveground biomass in 2017 and remained relatively stable throughout the winter season. In addition, night-time CO<sub>2</sub> fluxes were best correlated with soil temperature and WFPS ( $r = 0.69$  and  $-0.53$ , both  $P < 0.001$ ; Table 1). Overall, night-time CO<sub>2</sub> fluxes were negatively and positively correlated with CH<sub>4</sub> and N<sub>2</sub>O emissions, respectively ( $r = -0.41$  and  $0.46$ , both  $P < 0.001$ ).

### 3.3 Seasonal variability of ecosystem C fluxes

During the growing season, ecosystem C fluxes were measured seven times in total (Fig. 3). Marked fluctuations in ecosystem C fluxes occurred in July 2017, which may have resulted from substantial variations in soil WFPS (Fig. 3). Ecosystem C fluxes were strongly correlated with soil WFPS ( $r = 0.44$  to  $0.67$ ,  $P < 0.05$ – $0.001$ ; Table 1), while only the seasonal pattern of  $R_{eco}$  had a positive relationship with soil temperature ( $r = 0.45$ ,  $P < 0.05$ ).




**Figure 3** Soil temperature (A), water-filled pore space (WFPS) (B) at a depth of 10 cm, net ecosystem carbon exchange of CO<sub>2</sub> (NEE, C), ecosystem respiration ( $R_{eco}$ , D) and gross ecosystem productivity (GEP, E) during the growing season from a temperate semi-improved

grassland exposed to different levels of ozone and N. Values represent mean±SEM ( $n = 4$ ).

Results from the linear mixed effects model at  $P \leq 0.05$  are shown.

Across the growing season, elevated ozone significantly decreased NEE ( $P = 0.017$ ), whereas no effects of N fertilization ( $P = 0.823$ ) or its interaction with elevated ozone ( $P = 0.161$ ) were detected (Fig. 3C). Elevated ozone resulted in a 15% reduction of seasonal mean NEE. By contrast,  $R_{eco}$  was significantly increased by N fertilization ( $P = 0.028$ ) but not by elevated ozone or their interaction ( $P = 0.315$  and  $0.155$ ; Fig. 3D). Irrespective of ozone exposure, N fertilization significantly increased mean  $R_{eco}$  by 11%. Unlike NEE and  $R_{eco}$ , none of elevated ozone, N fertilization, or their interaction affected GEP (Fig. 3E).



**Figure 4** Aboveground biomass (A) and root biomass (B) in a temperate semi-improved grassland exposed to different levels of ozone and N. Values represent mean±SEM ( $n = 4$ ). Results from the linear mixed effects model at  $P \leq 0.05$  are shown.

### 3.4 Aboveground and root biomass

Harvested aboveground biomass differed significantly between cut events ( $P < 0.001$ ; Fig. 4A). In the 2017 autumn harvest, aboveground biomass from the N fertilized treatment (mean:  $703 \pm 23$  g DM m<sup>-2</sup>) was higher than from the control treatment (mean:  $515 \pm 20$  g DM m<sup>-2</sup>,  $P < 0.001$ ), but there was no difference between N treatments at the 2018 spring harvest ( $P = 0.736$ ). Aboveground biomass was affected by elevated ozone in the 2018 spring harvest but not in the 2017 autumn harvest. In the 2018 spring harvest, aboveground biomass decreased from  $238 \pm 12$  g DM m<sup>-2</sup> in the ambient ozone treatment to  $197 \pm 13$  g DM m<sup>-2</sup> in the elevated ozone treatment ( $P = 0.03$ ). The average root biomass was  $1450$  g DM m<sup>-2</sup> and did not differ between either elevated ozone or N fertilization treatments (Fig. 4B).

### *3.5 Soil properties, net N-cycling rates and enzyme activities*

Towards the end of the experiment, neither elevated ozone or N fertilization nor their interaction affected NH<sub>4</sub><sup>+</sup> concentrations, DON or microbial biomass C (Table 2). Nitrate concentrations and microbial biomass N were higher in the elevated ozone plots than in the ambient ozone plots ( $P = 0.003$  and  $0.012$ , respectively), but there was no difference for both parameters between control and N-fertilized plots. A marginally significant interaction between ozone and N fertilizer was detected for DOC ( $P = 0.081$ ). Microbial biomass C:N ratio was lower in the elevated ozone plots than in the ambient ozone plots ( $P = 0.018$ ). Net N mineralization, which is often used as an index of plant available mineral N, was higher in the elevated ozone treatment than in the ambient ozone treatment ( $P = 0.056$ ) but did not differ between the control

and N fertilized treatments. Further, net nitrification was higher in the elevated ozone than the ambient ozone treatment ( $P = 0.011$ ).

Potential activities in two of studied soil extracellular enzymes were not affected by elevated ozone, N fertilization or their interaction (Table 2). Elevated ozone and N fertilization significantly increased cellobiohydrolase activity by 26% and 73%, respectively ( $P = 0.04$  and  $0.003$ ). Elevated ozone also marginally stimulated leucine aminopeptidase activity by 15% as compared to under ambient ozone ( $P = 0.069$ ).

#### **4. Discussion**

In the past decade, many conclusions about the impacts of elevated ozone have been drawn from croplands (Bhatia et al., 2011; Decock et al., 2012; Kou et al., 2015; Tang et al., 2015; Zheng et al., 2011), peatlands (Toet et al., 2017, 2011) and subalpine grasslands (Volk et al., 2011). The effects of elevated ozone and N fertilization alone or in combination on GHG emissions and ecosystem C fluxes have been rarely reported, especially in temperate grasslands (Sánchez-Martín et al., 2017; Wang et al., 2019). This is to our knowledge the first study to report how GHG emissions, coupled with ecosystem C fluxes, in response to elevated ozone and N fertilization alone or in combination from a temperate semi-improved grassland under fully open-air field conditions.



**Table 2** Soil properties, net N-cycling rates and potential enzyme activities measured in May 2018 from a temperate semi-improved grassland exposed to different levels of ozone and N. Values represent mean $\pm$ SEM ( $n = 4$ ).

|   | Ambient ozone                 |                              | Elevated ozone               |                              |
|---|-------------------------------|------------------------------|------------------------------|------------------------------|
|   | Control                       | + Nitrogen                   | Control                      | + Nitrogen                   |
| NH <sub>4</sub> <sup>+</sup> (mg N kg <sup>-1</sup> ) | 4.48 $\pm$ 0.54               | 6.25 $\pm$ 0.54              | 5.30 $\pm$ 0.27              | 5.12 $\pm$ 0.90              |
| NO <sub>3</sub> <sup>-</sup> (mg N kg <sup>-1</sup> ) | 0.43 $\pm$ 0.03 <sup>B*</sup> | 0.46 $\pm$ 0.21 <sup>B</sup> | 1.28 $\pm$ 0.26 <sup>A</sup> | 1.01 $\pm$ 0.50 <sup>A</sup> |
| DOC (mg C kg <sup>-1</sup> )                          | 336 $\pm$ 13 <sup>†</sup>     | 295 $\pm$ 14                 | 314 $\pm$ 11                 | 318 $\pm$ 8                  |
| DON (mg N kg <sup>-1</sup> )                          | 59.7 $\pm$ 1.8                | 54.6 $\pm$ 3.1               | 55.6 $\pm$ 2.4               | 57.1 $\pm$ 1.0               |
| Microbial biomass                                     |                               |                              |                              |                              |
| C (mg C kg <sup>-1</sup> )                            | 2445 $\pm$ 67                 | 2371 $\pm$ 103               | 2548 $\pm$ 172               | 2461 $\pm$ 73                |
| Microbial biomass                                     |                               |                              |                              |                              |
| N (mg N kg <sup>-1</sup> )                            | 227 $\pm$ 3 <sup>B</sup>      | 229 $\pm$ 15 <sup>B</sup>    | 259 $\pm$ 24 <sup>A</sup>    | 253 $\pm$ 9 <sup>A</sup>     |
| Microbial biomass                                     |                               |                              |                              |                              |
| C:N   | 10.8 $\pm$ 0.3 <sup>A</sup>   | 10.4 $\pm$ 0.3 <sup>A</sup>  | 9.9 $\pm$ 0.3 <sup>B</sup>   | 9.7 $\pm$ 0.1 <sup>B</sup>   |
| Net N mineralization                                  |                               |                              |                              |                              |
| (mg N kg <sup>-1</sup> day <sup>-1</sup> )            | -0.03 $\pm$ 0.11 <sup>‡</sup> | -0.16 $\pm$ 0.09             | 0.22 $\pm$ 0.15              | 0.13 $\pm$ 0.15              |
| Net nitrification                                     |                               |                              |                              |                              |
| (mg N kg <sup>-1</sup> day <sup>-1</sup> )            | 0.04 $\pm$ 0.04 <sup>B</sup>  | 0.06 $\pm$ 0.09 <sup>B</sup> | 0.33 $\pm$ 0.06 <sup>A</sup> | 0.19 $\pm$ 0.08 <sup>A</sup> |
| $\beta$ -glucosidase                                  |                               |                              |                              |                              |
| (nmol g <sup>-1</sup> h <sup>-1</sup> )               | 126 $\pm$ 24                  | 121 $\pm$ 15                 | 113 $\pm$ 2                  | 132 $\pm$ 16                 |
| Cellobiohydrolase                                     |                               |                              |                              |                              |
| (nmol g <sup>-1</sup> h <sup>-1</sup> )               | 16.7 $\pm$ 2.8 <sup>Bb§</sup> | 27.1 $\pm$ 5.7 <sup>Ba</sup> | 19.7 $\pm$ 3.0 <sup>Ab</sup> | 36.0 $\pm$ 3.5 <sup>Aa</sup> |
| N-acetyl-glucosaminidase                              |                               |                              |                              |                              |
| (nmol g <sup>-1</sup> h <sup>-1</sup> )               | 83.6 $\pm$ 6.2                | 94.3 $\pm$ 7.5               | 83.5 $\pm$ 7.8               | 84.3 $\pm$ 9.9               |
| Leucine aminopeptidase                                |                               |                              |                              |                              |
| (nmol g <sup>-1</sup> h <sup>-1</sup> )               | 42.7 $\pm$ 3.4 <sup>‡</sup>   | 37.8 $\pm$ 1.7               | 45.1 $\pm$ 2.3               | 47.1 $\pm$ 3.8               |

\* Values with different uppercase letters indicate significant differences between ambient and elevated ozone treatments (linear mixed effects model with the Tukey HSD test at  $P \leq 0.05$ )

<sup>†</sup> Marginally significant interaction between ozone and N fertilized treatments (linear mixed effects model at  $P \leq 0.09$ )

<sup>‡</sup> Marginally significant effect of elevated ozone (linear mixed effects model at  $P \leq 0.09$ )

§ Values with different lowercase letters indicate significant differences between control and N-fertilized treatments (linear mixed effects model with the Tukey HSD test at  $P \leq 0.05$ )

#### *4.1 Response of CH<sub>4</sub> fluxes to elevated ozone and N fertilization*

The fluxes of CH<sub>4</sub> from our studied grasslands were generally negative throughout the experimental period, which agrees with the argument that temperate grassland soils generally act as an atmospheric CH<sub>4</sub> sink (Hötnagl et al., 2018; Smith et al., 2000). We found that elevated ozone tended to decrease the magnitude of CH<sub>4</sub> uptake (a slightly significant effect) compared to the ambient plots during the growing season but not the winter period, suggesting a negative impact on the CH<sub>4</sub> uptake capacity. This is unlikely to be related to a direct effect of ozone on soil methanogens and methanotrophs since ozone does not penetrate far into the soil (Blum and Tingey, 1977; Toet et al., 2009). The reduced CH<sub>4</sub> uptake during the growing season could be possibly due to altered methanotroph activity at elevated ozone. Across the one-year period CH<sub>4</sub> emissions were not affected by elevated ozone, which agrees with findings of several studies under OTC conditions (Kanerva et al., 2007; Sánchez-Martín et al., 2017). For example, studies in an annual Mediterranean grassland reveal that ozone exposure for a short-term period (49 days) did not affect CH<sub>4</sub> emissions (Sánchez-Martín et al., 2017). Similarly, the three-year field study of Kanerva et al. (2007) reported no overall effect of ozone exposure on CH<sub>4</sub> fluxes from ground-planted meadow mesocosms. By contrast, studies in rice fields under either OTC or O<sub>3</sub>-FACE conditions show that elevated ozone can decrease CH<sub>4</sub> emissions during the rice growing season (Bhatia et al., 2011; Tang et al., 2015; Zhang et al., 2016). Either

a negative or transient effect of elevated ozone on CH<sub>4</sub> emissions from peatlands is also reported (Mörsky et al., 2008; Toet et al., 2011).

Although it has been suggested that N addition decreases CH<sub>4</sub> uptake in upland ecosystems (Liu and Greaver, 2009), our results show that neither N fertilization nor its interaction with elevated ozone affected CH<sub>4</sub> uptake. This is supported by the findings of a meta-analysis where they found that soil CH<sub>4</sub> uptake in natural or with short-term N fertilization sites showed a non-significant change when N is added (Aronson and Helliker, 2010). Thus, the lack of CH<sub>4</sub> uptake in response to N addition could be attributed to a one-time application of N fertilizer into this semi-improved grassland. On the other hand, our result contradicts with their finding that N addition effects on soil CH<sub>4</sub> uptake may be switched from stimulation to inhibition when N addition exceeds a threshold value of 100 kg N ha<sup>-1</sup> yr<sup>-1</sup> (Aronson and Helliker, 2010). This implies that a shift from stimulation to inhibition of soil CH<sub>4</sub> uptake might be related to not only the amount of N added but also the inherent soil N status of the studied system. We are aware of only one previous field study where they examined the interaction effect between elevated ozone and N fertilization on GHG emissions from a simplified annual grassland (Sánchez-Martín et al., 2017). Consistent with our findings, the response of CH<sub>4</sub> emission to ozone exposure from their experiment site was not N fertilization dependent. While we found either elevated ozone or N fertilization may have a negligible effect on soil CH<sub>4</sub> uptake in this temperate grassland, future multiple-year studies are needed to reveal the inter-annual variability and its underlying mechanisms.

#### 4.2 Response of $N_2O$ fluxes to elevated ozone and N fertilization

This semi-improved grassland acted as a source of  $N_2O$ , although negative  $N_2O$  fluxes were recorded occasionally, especially during the winter season (Fig. 2D). Our results showed that neither elevated ozone or N fertilization nor their interaction had an appreciable effect on  $N_2O$  emission over the experimental year. The lack of responsiveness of  $N_2O$  emissions to N addition suggests that a one-time application of 100 kg N ha<sup>-1</sup> (the first fertilizer N application to this grassland for >25 years) at the beginning of the experiment would not have been sufficient to produce significant  $N_2O$  emissions in this semi-improved grassland. Annual  $N_2O$  emissions from this study were much higher (0.18–0.34 kg N ha<sup>-1</sup> yr<sup>-1</sup>) than that (ca. 0.06 kg N ha<sup>-1</sup> yr<sup>-1</sup>) from the field study of Marsden et al. (2018) in the same grassland. This is likely due to the differences in microclimate between the two study sites. In their study, the large amount of N applied in the form of sheep urine (ca. 2000 kg N ha<sup>-1</sup> yr<sup>-1</sup>) had no marked effect on  $N_2O$  fluxes (Marsden et al., 2018), which partly supports our finding. Furthermore, N fertilization significantly increased aboveground biomass during the growing season, but had minor resultant effects on plant yield and soil characteristics at the end of the experimental year (Fig. 4A; Table 2). Therefore, these findings collectively point to the consumption of applied N fertilizer tightly coupled to plant uptake and microbial immobilization in this semi-improved grassland.

To date, the divergent responses of  $N_2O$  fluxes to elevated ozone have been reported. For example, some studies showed a decrease in  $N_2O$  emissions under elevated ozone (Bhatia et al.,

2011; Kou et al., 2015), other reported stimulated emissions (Sánchez-Martín et al., 2017) or no marked effects (Decock et al., 2012; Kanerva et al., 2007) compared to the ambient ozone concentrations. As soil nitrification and denitrification processes are major pathways of N<sub>2</sub>O emissions, changes in the substrate availability due to elevated ozone are expected to alter N<sub>2</sub>O emissions. We found that elevated ozone resulted in a significant decline of aboveground biomass at the 2018 spring harvest, which may, in turn, contribute to accelerating soil N-cycling (i.e., a build-up of soil nitrate, higher rates of net N mineralization and nitrification; Table 2). The unchanged N<sub>2</sub>O emissions but stimulated N-cycling processes could be explained by the low intensity of soil N<sub>2</sub>O emission and/or other pathways being altered which might be primarily responsible for N loss from this system. Indeed, in a soybean agroecosystem under fully open-air conditions, Decock et al. (2012) also demonstrated that elevated ozone stimulated soil N-cycling but had no effect on soil N<sub>2</sub>O emissions. In contrast, in an annual Mediterranean grassland, ozone exposure reduced the fertilization effect of N on plant growth and thereby resulted in a significant reduction of soil N<sub>2</sub>O emissions (Sánchez-Martín et al., 2017). Clearly, this was not the case for our study because of the absence of an interaction between elevated ozone and N fertilization on plant yield, especially during the growing season (Fig. 4). Nevertheless, it should be noted that the single N application in our study might counterbalance the negative impact of ozone exposure on plant growth during the growing season.

#### *4.3 Effects of elevated ozone and N fertilization on ecosystem C fluxes*

A growing number of studies have documented the detrimental effects of elevated ozone on grassland plant species, ranging from visible injury of leaves to substantially decreased productivity (Hayes et al., 2007; Volk et al., 2006). In our study, elevated ozone resulted in a small reduction (3.2% for the control plots) but a significant reduction of 17.1% in aboveground biomass during the growing and non-growing seasons, respectively (Fig. 4A). During the growing season, elevated ozone caused a significant reduction of NEE (14.9%), which is consistent with other studies in an annual temperate grassland (Calvete-Sogo et al., 2014; Sánchez-Martín et al., 2017) and peatland microcosms (Haapala et al., 2011). Because of the unresponsiveness of total aboveground biomass, the reduced NEE by elevated ozone at the canopy level is likely due to the changed photosynthetic and/or respiration rates (Andersen, 2003). We noted that these semi-improved grassland mesocosms had a higher growth rate, in terms of stimulated fluxes of trace gas and night-time CO<sub>2</sub> (Fig. 2C-E), under the O<sub>3</sub>-FACE field condition than *in situ* on the upland site (Marsden et al., 2018). Thus, our results support the view that plant communities with fast-growing rates (i.e., annual and newly established systems) would be more responsive to elevated ozone than these perennial and well established systems that are characterized by slower growth rates (Grantz et al., 2006; Grime, 2000). In contrast, we are aware of a transient effect of elevated ozone on NEE in peatlands (Haapala et al., 2011; Niemi et al., 2002; Rinnan et al., 2003). On a subalpine grassland, Volk et al. (2011) found no effect of elevated ozone on NEE measured during the third growing season. Taken together, our results and those of previous findings suggest that further long-term investigations are warranted to evaluate the inter-annual response of NEE to elevated ozone in grasslands.

Ecosystem respiration as the sum of the respiration from plants and heterotrophs was not affected by elevated ozone during the growing season (Figs. 2E and 3D). This is in agreement with findings from other studies in subalpine and annual grasslands (Calvete-Sogo et al., 2014; Volk et al., 2011). Similarly, studies from peatlands suggested no effect of elevated ozone on ecosystem respiration, especially during the first experimental year (Haapala et al., 2011; Kanerva et al., 2007; Toet et al., 2011). In our study, the unchanged ecosystem respiration might be due to the response of heterotrophic respiration to elevated ozone which was unresponsive or masked by the larger proportion of plant respiration (Volk et al., 2011). As expected, stimulation of plant productivity provides more substrate for plant and soil respiration, leading to increases in ecosystem respiration following N fertilization (Figs. 2E and 3D). The positive response of ecosystem respiration to N addition is consistent with that in a subalpine grassland (Volk et al., 2011). In contrast, the lack of N effects on ecosystem respiration in an annual grassland was attributed to the fact that the low N doses were only enough to meet plant N demand but had no effect on the yield or gas exchange rates (Calvete-Sogo et al., 2014). Since the assimilation processes was primarily N-limited, we expected to see the positive response of GEP to N fertilization in this semi-improved grassland. However, the differential responses of NEE and ecosystem respiration to elevated ozone and N fertilization resulted in non-significant changes in GEP, suggesting N fertilization counterbalanced the effect of elevated ozone on GEP in our studied grassland.

#### *4.4 Short-term effects of elevated ozone and N fertilization on belowground processes*

While the intact grassland mesocosms were exposed to ozone only for one growing season, our results indeed showed that soil N dynamics were altered at the end of the experimental year. Increased soil mineral N availability under elevated ozone may be due to the significant reduction of aboveground biomass during the non-growing season, which may have contributed to reduced plant N uptake. On the other hand, we found that the activities of cellobiohydrolase and leucine aminopeptidase were higher in the elevated ozone plots than in the ambient plots, suggesting that elevated ozone stimulates microbes in the soil to produce enzymes that degrade cellulose and peptide-containing moieties. Parallel to the marginal increase of net N mineralization, our results suggest an increased soil N mineralization by elevated ozone in this grassland soil. Under elevated ozone, the unchanged soil  $\text{NH}_4^+$  is likely related to increased microbial N immobilization, whereas stimulated soil net nitrification may have resulted in an accumulation of soil  $\text{NO}_3^-$ . Consistent with our findings, other studies in the  $\text{O}_3$ -FACE systems have reported the increased soil mineral N availability under elevated ozone in both soybean and wheat fields (He et al., 2014; Wu et al., 2016). Note that our soil sampling was carried out in early spring of 2018, which probably contributes to masking the possible N fertilization effect or its interaction with ozone, especially during the growing season. As soil mineral N availability is mainly regulated by plant uptake and microbial-mediated N transformation processes (Schimel and Bennett, 2004), our results collectively point to a positive feedback of soil N transformation in this semi-improved grassland to short-term of ozone exposure.

## **5. Conclusions**



To our knowledge, this study is the first to report the effects of elevated ozone on GHG and ecosystem C fluxes in temperate grasslands under fully open-air field conditions. Our results demonstrate that elevated ozone may reduce atmospheric CH<sub>4</sub> uptake and net C uptake during the growing season in this semi-improved grassland. Given that the reported responses of CH<sub>4</sub> emission and NEE to elevated ozone are still inconsistent and controversial yet, we speculate that the depressed CH<sub>4</sub> uptake under elevated ozone could be transient, which warrants further investigation. The lack of any interaction between elevated ozone and N fertilization is likely due to the one-time N fertilization at the beginning of experiment, which probably was not synchronized with the occurrence of ozone injury of the plants. Further, our results suggest that short-term of ozone treatment may have contributed to accelerated soil N cycling in this grassland. Further investigations are warranted to examine the long-term impact of elevated ozone on ecosystem production and function.

### **Acknowledgements**

This work was supported by the European Commission under Horizon 2020 for a Marie Skłodowska-Curie Actions COFUND Fellowship (663830-BU-048); and the Welsh Government and Higher Education Funding Council for Wales through the Sêr Cymru National Research Network for Low Carbon, Energy and Environment. We thank Aled Williams for technical support in running the O<sub>3</sub>-FACE facility.

### **Reference**

540 Ainsworth, E., Yendrek, C.R., Sitch, S., Collins, W.J., Emberson, L.D., 2012. The effects of  
541 tropospheric ozone on net primary productivity and implications for climate change. *Annu.*  
542 *Rev. Plant Biol.* 63, 637–61. doi:10.1146/annurev-arplant-042110-103829

543 Andersen, C.P., 2003. Source-sink balance and carbon allocation below ground in plants  
544 exposed to ozone. *New Phytol.* 157, 213–228. doi:10.1046/j.1469-8137.2003.00674.x

545 Aronson, A.E.L., Helliker, B.R., 2010. Methane flux in non-wetland soils in response to  
546 nitrogen addition : a meta-analysis Linked references are available on JSTOR for this  
547 article : Methane flux in non-wetland soils in response to nitrogen addition : a meta-  
548 analysis. *Ecology* 91, 3242–3251. doi:10.2307/20788157

549 Bates, D., Mächler, M., Bolker, B., Walker, S., 2014. Fitting linear mixed-effects models using  
550 lme4. *arXiv Prepr. arXiv1406.5823*.

551 Bhatia, A., Ghosh, A., Kumar, V., Tomer, R., Singh, S.D., Pathak, H., 2011. Effect of elevated  
552 tropospheric ozone on methane and nitrous oxide emission from rice soil in north India.  
553 *Agric. Ecosyst. Environ.* 144, 21–28. doi:10.1016/j.agee.2011.07.003

554 Blum, U., Tingey, D.T., 1977. A study of the potential ways in which ozone could reduce root  
555 growth and nodulation of soybean. *Atmos. Environ.* 11, 737–739. doi:10.1016/0004-  
556 6981(77)90182-2

557 Brookes, P.C., Landman, A., Pruden, G., Jenkinson, D.S., 1985. Chloroform fumigation and  
558 the release of soil nitrogen: A rapid direct extraction method to measure microbial  
559 biomass nitrogen in soil. *Soil Biol. Biochem.* 17, 837–842. doi:10.1016/0038-

560 0717(85)90144-0

561 Calvete-Sogo, H., Elvira, S., Sanz, J., González-Fernández, I., García-Gómez, H., Sánchez-  
562 Martín, L., Alonso, R., Bermejo-Bermejo, V., 2014. Current ozone levels threaten gross  
563 primary production and yield of Mediterranean annual pastures and nitrogen modulates  
564 the response. *Atmos. Environ.* 95, 197–206. doi:10.1016/j.atmosenv.2014.05.073

565 Crawley, M.J., 2012. *The R book*. John Wiley & Sons.

566 Defra, 2016. *Agriculture in the United Kingdom 2015*. Department for Environment, Food and  
567 Rural Affairs, London, UK.

568 Decock, C., Chung, H., Venterea, R., Gray, S.B., Leahey, A.D.B., Six, J., 2012. Elevated CO<sub>2</sub>  
569 and O<sub>3</sub> modify N turnover rates, but not N<sub>2</sub>O emissions in a soybean agroecosystem. *Soil*  
570 *Biol. Biochem.* 51, 104–114. doi:10.1016/j.soilbio.2012.04.015

571 DeForest, J.L., 2009. The influence of time, storage temperature, and substrate age on potential  
572 soil enzyme activity in acidic forest soils using MUB-linked substrates and l-DOPA. *Soil*  
573 *Biol. Biochem.* 41, 1180–1186. doi:10.1016/j.soilbio.2009.02.029

574 FAO, 1981. *FAO-UNESCO Soil Map of the World 1:5000000*. Volume V: Europe, UNESCO,  
575 Paris, France.

576 Fowler, D., Amann, M., Anderson, F., Ashmore, M., Cox, P., Depledge, M., Derwent, D.,  
577 Grennfelt, P., Hewitt, N., Hov, O., Jenkin, M., Kelly, F., Liss, P., Pilling, M., Pyle, J., J,  
578 S., Stevenson, D., 2008. *Ground-level Ozone in the 21st Century: Future Trends, Impacts*  
579 *and Policy Implications*. *R. Soc. Sci. Policy Rep.* 15(08).

580 Fuhrer, J., Val Martin, M., Mills, G., Heald, C.L., Harmens, H., Hayes, F., Sharps, K., Bender,  
 581 J., Ashmore, M.R., 2016. Current and future ozone risks to global terrestrial biodiversity  
 582 and ecosystem processes. *Ecol. Evol.* 6, 8785–8799. doi:10.1002/ece3.2568  
 583 Grantz, D.A., Gunn, S., Vu, H.B., 2006. O<sub>3</sub> impacts on plant development: A meta-analysis of  
 584 root/shoot allocation and growth. *Plant, Cell Environ.* 29, 1193–1209.  
 585 doi:10.1111/j.1365-3040.2006.01521.x  
 586 Grime, J.P., 2000. The Response of Two Contrasting Limestone Grasslands to Simulated  
 587 Climate Change. *Science* (80-. ). 289, 762–765. doi:10.1126/science.289.5480.762  
 588 Haapala, J.K., Mörsky, S.K., Rinnan, R., Saarnio, S., Martikainen, P.J., Holopainen, T., Silvola,  
 589 J., 2011. Long-term effects of ozone on CO<sub>2</sub> exchange in peatland microcosms. *Atmos.*  
 590 *Environ.* 45, 4002–4007. doi:10.1016/j.atmosenv.2011.04.057  
 591 Hart, S.C., Stark, J.M., Davidson, E.A., Firestone, M.K., 1994. Nitrogen mineralization,  
 592 immobilization, and nitrification. *Methods Soil Anal. Part 2. Microbiol. Biochem. Prop.*  
 593 985–1018. doi:10.2136/sssabookser5.2.c42  
 594 Hayes, F., Jones, M.L.M., Mills, G., Ashmore, M., 2007. Meta-analysis of the relative  
 595 sensitivity of semi-natural vegetation species to ozone. *Environ. Pollut.* 146, 754–762.  
 596 doi:10.1016/j.envpol.2006.06.011  
 597 He, Z., Xiong, J., Kent, A.D., Deng, Y., Xue, K., Wang, G., Wu, L., Van Nostrand, J.D., Zhou,  
 598 J., 2014. Distinct responses of soil microbial communities to elevated CO<sub>2</sub> and O<sub>3</sub> in a  
 599 soybean agro-ecosystem. *ISME J.* 8, 714–26. doi:10.1038/ismej.2013.177

600 Hörtnagl, L., Barthel, M., Buchmann, N., Eugster, W., Butterbach-Bahl, K., Díaz-Pinés, E.,  
 601 Zeeman, M., Klumpp, K., Kiese, R., Bahn, M., Hammerle, A., Lu, H., Ladreiter-Knauss,  
 602 T., Burri, S., Merbold, L., 2018. Greenhouse gas fluxes over managed grasslands in  
 603 Central Europe. *Glob. Chang. Biol.* 24, 1843–1872. doi:10.1111/gcb.14079  
 604 IPCC, 2007. Summary for Policymakers, in: Solomon, S., Qin, D., Manning, M., Chen, Z.,  
 605 Marquis, M., Averyt, K.B., M.Tignor, Miller, H.L. (Eds.), *Climate Change 2007: The*  
 606 *Physical Science Basis. Contribution of Working Group I to the Fourth Assessment*  
 607 *Report of the Intergovernmental Panel on Climate Change.* Cambridge University Press,  
 608 Cambridge, United Kingdom and New York, NY, USA.  
 609 Kanerva, T., Regina, K., Rämö, K., Ojanperä, K., Manninen, S., 2007. Fluxes of N<sub>2</sub>O, CH<sub>4</sub> and  
 610 CO<sub>2</sub> in a meadow ecosystem exposed to elevated ozone and carbon dioxide for three years.  
 611 *Environ. Pollut.* 145, 818–828. doi:10.1016/j.envpol.2006.03.055  
 612 Kou, T.J., Cheng, X.H., Zhu, J.G., Xie, Z.B., 2015. The influence of ozone pollution on CO<sub>2</sub>,  
 613 CH<sub>4</sub>, and N<sub>2</sub>O emissions from a Chinese subtropical rice-wheat rotation system under  
 614 free-air O<sub>3</sub> exposure. *Agric. Ecosyst. Environ.* 204, 72–81.  
 615 doi:10.1016/j.agee.2015.02.013  
 616 Linn, D.M., Doran, J.W., 1984. Aerobic and Anaerobic Microbial Populations in No-till and  
 617 Plowed Soils1. *Soil Sci. Soc. Am. J.* 48, 794–799.  
 618 doi:10.2136/sssaj1984.03615995004800040019x  
 619 Liu, L., Greaver, T.L., 2009. A review of nitrogen enrichment effects on three biogenic GHGs:

620 the CO<sub>2</sub> sink may be largely offset by stimulated N<sub>2</sub>O and CH<sub>4</sub> emission. *Ecol. Lett.* 12,  
621 1103–1117. doi:10.1111/j.1461-0248.2009.01351.x

622 Luyssaert, S., Inglima, I., Jung, M., Richardson, A.D., Reichstein, M., Papale, D., Piao, S.L.,  
623 Schulze, E.D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beer, C., Bernhofer,  
624 C., Black, K.G., Bonal, D., Bonnefond, J.M., Chambers, J., Ciais, P., Cook, B., Davis,  
625 K.J., Dolman, A.J., Gielen, B., Goulden, M., Grace, J., Granier, A., Grelle, A., Griffis, T.,  
626 Gr??nwald, T., Guidolotti, G., Hanson, P.J., Harding, R., Hollinger, D.Y., Hutyra, L.R.,  
627 Kolari, P., Kruijt, B., Kutsch, W., Lagergren, F., Laurila, T., Law, B.E., Le Maire, G.,  
628 Lindroth, A., Loustau, D., Malhi, Y., Mateus, J., Migliavacca, M., Misson, L.,  
629 Montagnani, L., Moncrieff, J., Moors, E., Munger, J.W., Nikinmaa, E., Ollinger, S. V.,  
630 Pita, G., Rebmann, C., Rouspard, O., Saigusa, N., Sanz, M.J., Seufert, G., Sierra, C., Smith,  
631 M.L., Tang, J., Valentini, R., Vesala, T., Janssens, I.A., 2007. CO<sub>2</sub> balance of boreal,  
632 temperate, and tropical forests derived from a global database. *Glob. Chang. Biol.* 13,  
633 2509–2537. doi:10.1111/j.1365-2486.2007.01439.x

634 Marsden, K.A., Holmberg, J.A., Jones, D.L., Chadwick, D.R., 2018. Sheep urine patch N<sub>2</sub>O  
635 emissions are lower from extensively-managed than intensively-managed grasslands.  
636 *Agric. Ecosyst. Environ.* 265, 264–274. doi:10.1016/j.agee.2018.06.025

637 Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., Gaye, T., Gregory, J.M., Kitoh,  
638 A., Knutti, R., Murphy, J.M., Noda, A., Raper, S.C.B., Watterson, I.G., Weaver, A.J.,  
639 Zhao, Z.C., 2007. Global climate projections, in: Solomon, S., Qin, D., Manning, M.,  
640 Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), IPCC, 2007:

641 Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to  
 642 the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.  
 643 Cambridge, UK, Cambridge University Press, pp. 747–846.

644 Mills, G., Harmens, H., Wagg, S., Sharps, K., Hayes, F., Fowler, D., Sutton, M., Davies, B.,  
 645 2016. Ozone impacts on vegetation in a nitrogen enriched and changing climate. *Environ.*  
 646 *Pollut.* 208, 898–908. doi:10.1016/j.envpol.2015.09.038

647 Mills, G., Hayes, F., Simpson, D., Emberson, L., Norris, D., Harmens, H., Büker, P., 2011.  
 648 Evidence of widespread effects of ozone on crops and (semi-)natural vegetation in Europe  
 649 (1990-2006) in relation to AOT40- and flux-based risk maps. *Glob. Chang. Biol.* 17, 592–  
 650 613. doi:10.1111/j.1365-2486.2010.02217.x

651 Miranda, K.M., Espey, M.G., Wink, D.A., 2001. A rapid, simple spectrophotometric method  
 652 for simultaneous detection of nitrate and nitrite. *Nitric Oxide - Biol. Chem.* 5, 62–71.  
 653 doi:10.1006/niox.2000.0319

654 Mörsky, S.K., Haapala, J.K., Rinnan, R., Tiiva, P., Saarnio, S., Silvola, J., Holopainen, T.,  
 655 Martikainen, P.J., 2008. Long-term ozone effects on vegetation, microbial community and  
 656 methane dynamics of boreal peatland microcosms in open-field conditions. *Glob. Chang.*  
 657 *Biol.* 14, 1891–1903. doi:10.1111/j.1365-2486.2008.01615.x

658 Mulvaney, R.L., 1996. Nitrogen—Inorganic Forms, in: Sparks, D.L. (Ed.), *Methods of Soil*  
 659 *Analysis Part 3—Chemical Methods*. Soil Science Society of America, American Society  
 660 of Agronomy, Madison, WI, pp. 1123–1184. doi:10.2136/sssabookser5.3.c38

661 Niemi, R., Martikainen, P.J., Silvola, J., Holopainen, T., 2002. Ozone effects on Sphagnum  
 662 mosses, carbon dioxide exchange and methane emission in boreal peatland microcosms.  
 663 Sci. Total Environ. 289, 1–12. doi:10.1016/S0048-9697(01)01012-9  
 664 Niu, S., Wu, M., Han, Y., Xia, J., Zhang, Z., Yang, H., Wan, S., 2010. Nitrogen effects on net  
 665 ecosystem carbon exchange in a temperate steppe. Glob. Chang. Biol. 16, 144–155.  
 666 doi:10.1111/j.1365-2486.2009.01894.x  
 667 R Development Core Team, 2015. R: A language and environment for statistical computing.  
 668 Vienna, Austria.  
 669 Rinnan, R., Impiö, M., Silvola, J., Holopainen, T., Martikainen, P.J., 2003. Carbon dioxide and  
 670 methane fluxes in boreal peatland microcosms with different vegetation cover - Effects of  
 671 ozone or ultraviolet-B exposure. Oecologia 137, 475–483. doi:10.1007/s00442-003-  
 672 1366-5  
 673 Rodwell, J.S., 1992. British plant communities. Volume 3. Grasslands and montane  
 674 communities, Cambridge University Press, Cambridge, UK. Cambridge University Press,  
 675 Cambridge, UK.  
 676 Saiya-Cork, K., Sinsabaugh, R., Zak, D., 2002. The effects of long term nitrogen deposition  
 677 on extracellular enzyme activity in an Acer saccharum forest soil. Soil Biol. Biochem. 34,  
 678 1309–1315. doi:10.1016/S0038-0717(02)00074-3  
 679 Sánchez-Martín, L., Bermejo-Bermejo, V., García-Torres, L., Alonso, R., de la Cruz, A.,  
 680 Calvete-Sogo, H., Vallejo, A., 2017. Nitrogen soil emissions and belowground plant



681 processes in Mediterranean annual pastures are altered by ozone exposure and N-inputs.  
 682 Atmos. Environ. 165, 12–22. doi:10.1016/j.atmosenv.2017.06.030

683 Schimel, J.P., Bennett, J., 2004. Nitrogen mineralization: Challenges of a changing paradigm.  
 684 Ecology 85, 591–602. doi:10.1890/03-8002

685 Sicard, P., Anav, A., De Marco, A., Paoletti, E., 2017. Projected global ground-level ozone  
 686 impacts on vegetation under different emission and climate scenarios. Atmos. Chem. Phys.  
 687 17, 12177–12196. doi:10.5194/acp-17-12177-2017

688 Sitch, S., Cox, P.M., Collins, W.J., Huntingford, C., 2007. Indirect radiative forcing of climate  
 689 change through ozone effects on the land-carbon sink. Nature 448, 791–794.  
 690 doi:10.1038/nature06059

691 Smith, K.A., Dobbie, K.E., Ball, B.C., Bakken, L.R., Sitaula, B.K., Hansen, S., Brumme, R.,  
 692 Borken, W., Christensen, S., Priemé, A., Fowler, D., Macdonald, J.A., Skiba, U.,  
 693 Klemetsson, L., Kasimir-Klemetsson, A., Degórska, A., Orlanski, P., 2000. Oxidation  
 694 of atmospheric methane in Northern European soils, comparison with other ecosystems,  
 695 and uncertainties in the global terrestrial sink. Glob. Chang. Biol. 6, 791–803.  
 696 doi:10.1046/j.1365-2486.2000.00356.x

697 Tang, H., Liu, G., Zhu, J., Kobayashi, K., 2015. Effects of elevated ozone concentration on CH<sub>4</sub>  
 698 and N<sub>2</sub>O emission from paddy soil under fully open-air field conditions. Glob. Chang.  
 699 Biol. 21, 1727–1736. doi:10.1111/gcb.12810

700 Toet, S., Ineson, P., Peacock, S., Ashmore, M., 2011. Elevated ozone reduces methane

701 emissions from peatland mesocosms. *Glob. Chang. Biol.* 17, 288–296.  
 702 doi:10.1111/j.1365-2486.2010.02267.x

703 Toet, S., Oliver, V., Ineson, P., McLoughlin, S., Helgason, T., Peacock, S., Stott, A.W., Barnes,  
 704 J., Ashmore, M., 2017. How does elevated ozone reduce methane emissions from  
 705 peatlands? *Sci. Total Environ.* 579, 60–71. doi:10.1016/j.scitotenv.2016.10.188

706 Toet, S., Subke, J.A., D’Haese, D., Ashmore, M.R., Emberson, L.D., Crossman, Z., Evershed,  
 707 R.P., Barnes, J.D., Ineson, P., 2009. A new stable isotope approach identifies the fate of  
 708 ozone in plant-soil systems. *New Phytol.* 182, 85–90. doi:10.1111/j.1469-  
 709 8137.2009.02780.x

710 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil  
 711 microbial biomass C. *Soil Biol. Biochem.* 19, 703–707. doi:10.1016/0038-  
 712 0717(87)90052-6

713 Volk, M., Bungener, P., Contat, F., Montani, M., Fuhrer, J., 2006. Grassland yield declined by  
 714 a quarter in 5 years of free-air ozone fumigation. *Glob. Chang. Biol.* 12, 74–83.  
 715 doi:10.1111/j.1365-2486.2005.01083.x

716 Volk, M., Obrist, D., Novak, K., Giger, R., Bassin, S., Fuhrer, J., 2011. Subalpine grassland  
 717 carbon dioxide fluxes indicate substantial carbon losses under increased nitrogen  
 718 deposition, but not at elevated ozone concentration. *Glob. Chang. Biol.* 17, 366–376.  
 719 doi:10.1111/j.1365-2486.2010.02228.x

720 Volk, M., Wolff, V., Bassin, S., Ammann, C., Fuhrer, J., 2014. High tolerance of subalpine

721 grassland to long-term ozone exposure is independent of N input and climatic drivers.  
 722 Environ. Pollut. 189, 161–168. doi:10.1016/j.envpol.2014.02.032

723 Wang, J., Hayes, F., Turner, R., Chadwick, D.R., Mills, G., Jones, D.L., 2019. Effects of four  
 724 years of elevated ozone on microbial biomass and extracellular enzyme activities in a  
 725 semi-natural grassland. Sci. Total Environ. 660, 260–268.  
 726 doi:10.1016/j.scitotenv.2019.01.040

727 Williamson, J.L., Mills, G., Hayes, F., Jones, T., Freeman, C., 2016. How do increasing  
 728 background concentrations of tropospheric ozone affect peatland plant growth and carbon  
 729 gas exchange? Atmos. Environ. 127, 133–138. doi:10.1016/j.atmosenv.2015.12.004

730 Wu, H., Li, Q., Lu, C., Zhang, L., Zhu, J., Dijkstra, F.A., Yu, Q., 2016. Elevated ozone effects  
 731 on soil nitrogen cycling differ among wheat cultivars. Appl. Soil Ecol. 108, 187–194.  
 732 doi:10.1016/j.apsoil.2016.08.015

733 Wu, J., Joergensen, R., Pommerening, B., Chaussod, R., Brookes, P.C., 1990. Measurement of  
 734 soil microbial biomass C by fumigation-extraction—an automated procedure. Soil Biol.  
 735 Biochem. 22, 1167–1169. doi:10.1016/0038-0717(90)90046-3

736 Yendrek, C.R., Leisner, C.P., Ainsworth, E.A., 2013. Chronic ozone exacerbates the reduction  
 737 in photosynthesis and acceleration of senescence caused by limited N availability in  
 738 *Nicotiana sylvestris*. Glob. Chang. Biol. 19, 3155–3166. doi:10.1111/gcb.12237

739 Zhang, J., Tang, H., Zhu, J., Lin, X., Feng, Y., 2016. Divergent responses of methanogenic  
 740 archaeal communities in two rice cultivars to elevated ground-level O<sub>3</sub>. Environ. Pollut.

741 213, 127–134. doi:10.1016/j.envpol.2016.01.062

742 Zheng, F., Wang, X., Lu, F., Hou, P., Zhang, W., Duan, X., Zhou, X., Ai, Y., Zheng, H., Ouyang,

743 Z., Feng, Z., 2011. Effects of elevated ozone concentration on methane emission from a

744 rice paddy in Yangtze River Delta, China. *Glob. Chang. Biol.* 17, 898–910.

745 doi:10.1111/j.1365-2486.2010.02258.x

746